


Water and Children's Health

Country	% of children under 5 with diarrhoea in the two weeks prior to the survey*	% with diarrhoea that received ORS packet	% with diarrhoea that received ORS or SSS	% with diarrhoea that received increased fluids plus continued feeding	Source	Year
Afghanistan	20.0	33.0			Multiple Indicator Cluster Survey	1997
Azerbaijan	21.7	10.1		26.5	Multiple Indicator Cluster Survey	2000
Bangladesh	6.1	61.4	73.6		Demographic and Health Survey	2000
Cambodia	18.9	17.8			Demographic and Health Survey	2000
China		11.3		28.8	National ORT Use Survey	1994
India	19.2	26.8			Demographic and Health Survey	1999
Indonesia	10.4	47.7		28.1	Demographic and Health Survey	1997
Kazakhstan	13.4	32.0		19.5	Demographic and Health Survey	1999
Kyrgyzstan	17.6	40.4	43.9	13.4	Demographic and Health Survey	1997
Lao PDR	6.2	33.9		20.1	Multiple Indicator Cluster Survey	2000
Maldives	7.9	51.2			Multiple Indicator Cluster Survey	1996
Mongolia	8.0	56.0		32.0	Multiple Indicator Cluster Survey	2000
Myanmar				23.9	Multiple Indicator Cluster Survey	1995
Nepal	27.5	25.9		11.2	Demographic and Health Survey	1996
Pakistan	26.0	46.0		19.0	Multiple Indicator Cluster Survey	1995
Papua New Guinea	16.5	16.7			Demographic and Health Survey	1996
Philippines	7.4	43.4		28.3	Demographic and Health Survey	1998
Sri Lanka	5.0	34.4			Demographic and Health Survey	1993
Tajikistan	20.8	35.3		19.9	Multiple Indicator Cluster Survey	2000
Thailand		73.3			Thailand Provincial Health Survey	1995
Turkmenistan	6.7	57.7		31.0	Multiple Indicator Cluster Survey	1995
Uzbekistan	5.3	32.0		18.5	Multiple Indicator Cluster Survey	2000
Viet Nam	11.3	11.2		20.2	Multiple Indicator Cluster Survey	2000

* The prevalence of diarrhoea may vary by season. Country surveys were administered at different times, rendering the diarrhoea prevalence data incomparable between countries.

Source: UNICEF

Water from Transboundary Rivers as Share of Annual Water Resources



Source: World Resources Institute (<http://www.earthtrends.wri.org>)